Laser assisted proton collision on light nuclei at moderate energies

I. F. Barna and S. Varró
Wigner Research Centre of the Hungarian Academy of Sciences,
Konkoly Thege út 29 - 33, 1121 Budapest, Hungary
and
ELI-HU Nonprofit Kft.,
Dugonics Tér 13, 6720 Szeged, Hungary
(Dated: June 25, 2014)

We present analytic angular differential cross section model for laser assisted proton nucleon scattering on a Woods-Saxon optical potential where the nth-order photon absorption is taken into account simultaneously. As a physical example we calculate cross sections for proton - ^{12}C collision at 49 MeV in the laboratory frame where the laser intensity is in the range of $10^7 - 10^{21}$ W/cm^2 at optical frequencies. The upper intensity limit is slightly below the relativistic regime.

PACS numbers: 24.10.Ht,25.40.Ep,32.80.Wr

arXiv:1406.6310v1 [nucl-th] 24 Jun 2014

I. INTRODUCTION

Nowadays optical laser intensities exceeded the 10^{22} W/cm^2 limit where radiation effects dominate the electron dynamics. In the field of laser-matter interaction a large number of non-linear response of atoms, molecules and plasmas can be investigated both theoretically and experimentally. Such interesting high-field phenomena are high harmonic generations, or plasma-based laser-electron acceleration. This field intensities open the door to high field quantum electrodynamics phenomena like vacuum-polarization effects of pair production [1]. In most of the presented studies the dynamics of the participating electrons are investigated. Numerous surveys on laser assisted electron collisions are available as well [2]. However, there are only few nuclear photo-excitation investigations done where some low-lying first excited states of medium of heavy elements are populated with the help of x-ray free electron laser pulses [3]. Nuclear excitation by atomic electron re-scattering in a laser field was investigated as well [4]. Various additional concepts are under consideration for photo-nuclear reactions by laser-driven gamma beams [5].

To our knowledge there are no publications available where laser assisted proton nucleus collisions (or radiative proton-nucleus scattering) were investigated. This is the goal of our recent paper. We consider the full optical potential of Woods-Saxon(WS) type [7] with the proper parametrization for moderate energy proton - ^{12}C collision [8]. The optical potential formalism has been a very successful method to study the single particle spectra of nucleus in the last four decades. Detailed description of the validity of this formalism can be found in a nuclear physics textbooks or in monographs like [9, 10].

The nuclear physics community recently managed to evaluate the closed analytic form of the Fourier transformed WS interaction [11] which is a great success. We incorporate these results into a first Born approximation scattering cross section formula where the initial and final proton wave functions are Volkov waves and the induced photon emission and absorption processes are taken into account up to arbitrary orders [12–18]. We end our study with a physical example where 49 MeV protons are scattered on ^{12}C nuclei in the field of a Titan sapphire laser.

II. THEORY

In the following we summarize our applied non-relativistic quantum mechanical description. The laser field is handled in the classical way via the minimal coupling. The laser beam is taken to be linearly polarized and the dipole approximation is used. If the dimensionless intensity parameter (or the normalised vector potential) $a_0 = 8.55 \cdot 10^{-10}\sqrt{I(\frac{W}{cm^2})}\lambda(\mu m)$ of the laser field is smaller than unity the non-relativistic description in dipole approximation is valid. For 800 nm laser wavelength this means a critical intensity of $I = 2.13 \cdot 10^{18}$W/cm^2. In case of protons a_0 is replaced by $a_p = ((m_p/m_e)^{-1})a_0$, where the proton to electron mass ratio is (m_p/m_e)=1836 [19]). Accordingly, for 800 nm wavelength the critical intensity for protons is $I_{crit} = 3.91 \cdot 10^{21}$ W/cm^2. Additionally, we consider moderate proton kinetic energy, not so much above the Coulomb barrier and neglect the interchange term between the proton projectile an the target carbon protons. This proton exchange effect could be included in the presented model with the help of Woods-Saxon potentials of non-local type [20] but not in the scope of the recent study.

To describe the scattering process of a proton on a nucleus in a spherically symmetric field the following Schrödinger

equation has to be solved,

$$\left[\frac{1}{2m}\left(\mathbf{p}-\frac{e}{c}\mathbf{A}\right)^2+U(r)\right]\Psi=i\hbar\frac{\partial\Psi}{\partial t}, \tag{1}$$

where $\mathbf{p}, \Psi$ are the momentum operator and the wave function of the proton, $\mathbf{A}(t)=A_0\epsilon cos(\omega t)$ is the vector potential of the external laser field with unit polarisation vector ϵ and $U(r)$ is the scattering potential. We search the solution in the following form of

$$\Psi=\varphi_{p_i}+\Psi_c, \tag{2}$$

where

$$\varphi_p(\mathbf{r},t)=\frac{1}{(2\pi\hbar)^{3/2}}e^{\frac{i}{\hbar}(\mathbf{pr}-Et)}exp\left[-\frac{i}{\hbar}\frac{p^2t}{2m}-\int_0^t dt'\left(\frac{e\mathbf{pA}(t')}{mc}+\frac{1}{2m}\left(\frac{eA(t')}{c}\right)^2\right)\right] \tag{3}$$

is the Volkov wave function of the scattering proton. We note at this point that the A^2 term drops out from the transition amplitude, because it depends only on time (and does not depend on the proton's momentum). The correction term is

$$\Psi_c=\int d^3p a_{\mathbf{p}}(t)\varphi_p, \tag{4}$$

where $a_{\mathbf{p}}(t)$ is the scattering amplitude. Calculating the transition matrix element between the final and the initial states in the first Born approximation for $a_{\mathbf{p}_f}(t)$ at $t\to\infty$ we get

$$T_{fi}=\langle\varphi_f|U|\Psi_i\rangle=-\frac{U(\mathbf{q})}{(2\pi\hbar)^3}\sum_{n=-\infty}^{\infty}2\pi i\delta\left(\frac{p_f^2-p_i^2}{2m}+n\hbar\omega\right)J_n(z), \tag{5}$$

where $U(\mathbf{q})$ is the Fourier transformed of the scattering potential with the momentum transfer of $\mathbf{q}\equiv\mathbf{p}_i-\mathbf{p}_f$ where $\mathbf{p}_i$ is the initial and $\mathbf{p}_f$ is the final proton momenta. The absolute value is $q=\sqrt{p_i^2+p_f^2-2p_ip_fcos(\theta_{p_i,p_f})}$. In our case, for 49 MeV energy protons absorbing optical photons the following approximation is valid $q\approx 2p_i|sin(\theta/2)|$.

The Dirac delta describes photon absorptions ($n<0$) and emissions ($n>0$). $J_n(z)$ is the Bessel function with the argument of

$$z\equiv\frac{m_e}{m_p}a_0(\hat{\mathbf{q}}\epsilon)\frac{2p_i}{\hbar k_0}|sin(\theta/2)| \tag{6}$$

where m_e, m_p are the electron and proton masses, a_0 is the dimensionless intensity parameter, (given above), $\hat{\mathbf{q}}$ and ϵ are the unit vectors of the momentum transfer and the laser polarisation direction. It can be shown with geometrical means that for low energy photons where ($E_{ph}<E_{p^+}$) the angle in the scalar product of $\hat{\mathbf{q}}\epsilon\equiv cos\chi$ is $\chi=\pi/2-\theta/2$ where θ is the scattering angle of the proton varying from 0 to π.
From $\frac{p_i}{\hbar k_0}=\sqrt{\frac{m_p}{m_e}}\sqrt{\frac{2m_ec^2E_i}{\hbar^2\omega_0^2}}$ collecting the constants together the final formula for z reads

$$z=\frac{1.4166\times10^{-3}}{\hbar\omega_0}\sqrt{\frac{E_p}{1836}}\sqrt{I}\times cos(\chi)\times|sin(\theta/2)|, \tag{7}$$

where the laser energy $\hbar\omega_0$ is measured in eV, the proton energy E_p in MeV and the laser intensity I in W/cm^2.

The final differential cross section formula for the laser associated collision with simultaneous nth-order photon absorption and emission processes is

$$\frac{d\sigma^{(n)}}{d\Omega}=\frac{p_f}{p_i}J_n^2(z)\frac{d\sigma_B}{d\Omega}. \tag{8}$$

The $\frac{d\sigma_B}{d\Omega}=\left(\frac{m}{2\pi\hbar^2}\right)^2|U(\mathbf{q})|^2$ is the usual Born cross section for the scattering on the potential U(r) alone (without the laser field). The expression Eq. (8) was calculated with different authors using different methods [12–18].

The scattering interaction $U(r)$ is the sum of the Coulomb potential of a uniform charged sphere [6] and a short range optical Woods-Saxon potential [7]

$$U(r) = V_c(r) + V_{ws}(r) + i[W(r) + W_s(r)] + V_{ls}(r)\mathbf{l}\cdot\sigma \tag{9}$$

where the Coulomb term is

$$\begin{aligned} V_c &= \frac{Z_p Z_t e^2}{2R_0}\left(3 - \frac{r^2}{R_c^2}\right) \qquad r < R_c \\ V_c &= \frac{Z_p Z_t e^2}{r} \qquad r \geq R_c \end{aligned} \tag{10}$$

where $R_c = r_0 A_t^{1/3}$ is the target radius calculated from the mass number of the target with $r_0 = 1.25fm$. Z_p, Z_t are the charge of the projectile and the target and e is the elementary charge. This kind of regularised Coulomb potential helps us to avoid singular cross sections and routinely used in nuclear physics.

The short range nuclear part is given via

$$\begin{aligned} V(r) &= -V_r f_{ws}(r, R_0, a_0) \\ W(r) &= -V_v f_{ws}(r, R_s, a_s) \\ W_s(r) &= -W_s(-4a_s) f'_{ws}(r, R_s, a_s) \\ V_{ls}(r) &= -(V_{so} + iW_{so})(-2) g_{ws}(r, R_{so}, a_{so}) \\ f_{ws}(r, R, a) &= \frac{1}{1 + exp\left(\frac{r-R}{a}\right)} \\ f'_{ws}(r, R, a) &= \frac{d}{dr} f_{ws}(r, R, a) \\ g_{ws}(r, R, a) &= f'_{ws}(r, R, a)/r. \end{aligned} \tag{11}$$

The constants V_r, W_v, V_{so} and W_{so} are the strength parameters, and $a_{0,s,so}, R_{0,s,so}$ are the diffuseness and the radius parameters given for large number of nuclei. As we will see at moderate collisions energies the complex tems become zero. In the last part of the present paper we will use the numerical parameters of [8] for proton carbon-collision.

According to the work of Hlophe [11] the complete analytic form of the Fourier transform of the Woods-Saxon potential is available

$$V_s(q) = \frac{V_r}{\pi^2}\left\{\frac{\pi a_0 e^{-\pi a_0 q}}{q(1 - e^{-2\pi a_0 q})^2}\left[R_0(1 - e^{-2\pi a_0 q})cos(qR_0) - \pi a_0(1 + e^{-2\pi a_0 q})sin(qR_0)\right] - a_0^3 e^{-\frac{R_0}{a_0}}\left[\frac{1}{(1 + a_0^2 q^2)^2} - \frac{2e^{-\frac{R_0}{a_0}}}{(4 + a_0^2 q^2)^2}\right]\right\}. \tag{12}$$

For the $W(q)$ imaginary term, the same expression was derived with W_v, a_s, R_s instead of V_r, a_0 and R_0. The surface term $W_s(r)$ gives the following formula in the momentum space:

$$W_s(q) = -4a_s\frac{W_s}{\pi^2}\left\{\frac{\pi a_s e^{-\pi a_s q}}{(1 - e^{-2\pi a_s q})^2}\left[(\pi a_s(1 + e^{-2\pi a_s q}) - \frac{1}{q}(1 - e^{-2\pi a_s q}))cos(qR_s) + R_s(1 - e^{-2\pi a_s q})sin(qR_s)\right] + a_s^2 e^{-\frac{R_s}{a_s}}\left[\frac{1}{(1 + a_s^2 q^2)^2} - \frac{4e^{-\frac{R_s}{a_s}}}{(4 + a_s^2 q^2)^2}\right]\right\}. \tag{13}$$

The transformed spin-orbit coupling term leads to

$$V_{ls}(q) = -\frac{a_{so}}{\pi^2}(V_{so} + iW_{so})\left\{\frac{2\pi e^{-\pi a_{so} q}}{1 - e^{-2\pi a_{so} q}} sin(qR_{so}) + e^{\frac{-R_{so}}{a_{so}}}\left(\frac{1}{1 + a_{so}^2 q^2} - \frac{2e^{-R_{so}}}{4 + a_{so}^2 q^2}\right)\right\}. \tag{14}$$

where the momentum transfer is defined as above $\mathbf{q} \equiv \mathbf{p_i} - \mathbf{p_f}$. The low energy transfer approximation formula $q \approx 2p_i|sin(\theta/2)|$ is valid.

The Fourier transform of the charged sphere Coulomb field is also far from being trivial

$$\begin{aligned} V_c(q) &= \frac{Z_p Z_t e^2}{2^{\frac{5}{6}}\sqrt{\pi} q^3}\left(-2 \cdot 3^{\frac{1}{3}} q \cos[2^{\frac{2}{3}} 3^{\frac{1}{3}} q] + 2^{\frac{1}{3}}(1 + 2 \cdot 2^{\frac{1}{3}} 3^{\frac{2}{3}} q^2)\sin[2^{\frac{2}{3}} 3^{\frac{1}{3}} q]\right) + \\ & 3 Z_p Z_t e^2 \sqrt{\frac{2}{\pi}}\left(\frac{i\pi|q|}{2q} - \mathrm{Ci}[2^{\frac{2}{3}} 3^{\frac{1}{3}} q] + \log(q) - \log|q| - i\mathrm{Si}[2^{\frac{2}{3}} 3^{\frac{1}{3}} q]\right) \end{aligned} \tag{15}$$

where Ci and Si are the cosine integral and the sine integral functions, respectively.



FIG. 1: The angular differential cross sections in the first Born approximation of the various Woods-Saxon potentials tems for 49 MeV elastic proton - ^{12}C scattering. The solid, dashed, and dotted lines are the contributions of Eqs. (12,13,14), respectively. Note, the different smoothness and different back scattering values of the different terms.

III. RESULTS

We applied the outlined method to 49 MeV proton - ^{12}C scattering. Table I contains the parameters of the used Woods-Saxon potential.

Name of the parameter	Numerical value
V_R	31.31 (MeV)
R_0	1.276 (fm)
a_0	0 .68 (fm)
W_s	5.98 (MeV)
r_s	0.890 (fm)
a_s	0.586 (fm)
V_{so}	2.79 (MeV)
R_{so}	0.716 (fm)
a_{so}	0 .222 (fm)

Table 1. The parameters of the applied potential for proton - ^{12}C collision at $E_i = 49$ MeV.
Note, that the complex part W_v and the the complex part of the spin-orbit term W_{so} are zero at this energy.

Figure 1 presents the angular differential cross section in the first Born approximation of the various Woods-Saxon potential terms for 49 MeV elastic proton - ^{12}C scattering. The different lines represent the different terms (12,13,14). The laboratory frame is used in the calculation. For a better transparency the contributions of the regularised Coulomb term is not presented. Our calculated total cross section of the elastic scattering is 201 mbarn which is consistent with the data of [8].

In our case the laser photon energy is $\hbar\omega_0 = 1.56$ eV, which means 800 nm wavelength and the proton energy is $E_i = 49$ MeV. With these values the argument of the Bessel function Eq. (7) becomes the following

$$z = 1.48 \times 10^{-3}\sqrt{I} \times cos(\chi) \times |sin(\theta/2)| = \mathcal{I} \times cos(\chi) \times |sin(\theta/2)|. \tag{16}$$

Figure 2 presents the angular differential cross section for $n = 0, 1, 2, 3$ photon absorptions for $\mathcal{I} = 1$ which means $I = 4.56 \times 10^7$ W/cm^2 intensity. The single, double and triple photon absorption total cross sections are 2.26, 1.1 and 0.71 mbarn, respectively.

Figure 3 shows the angular differential cross section for $n = 0, 1, 2$ photon absorptions for $\mathcal{I} = 100$ which means $I = 4.56 \times 10^{11}$ W/cm^2 intensity. Note, that the cross sections for single and double photon absorption are almost the same. The single photon absorption total cross section is 0.5 mbarn.



FIG. 2: The calculated angular differential cross sections from Eq. (8) for $I = 4.56 \times 10^7$ W/cm^2 laser field intensity ($\mathcal{I} = 1$). The thick solid, thin long dashed, thin short dashed and thin solid lines are for $n = 0, 1, 2, 3$ photon absorptions, respectively.



FIG. 3: The calculated angular differential cross sections from Eq. (8) for $I = 4.56 \times 10^{11}$ W/cm^2 laser field intensity ($\mathcal{I} = 100$). The thick solid, thin long dashed, thin short dashed are for $n = 0, 1, 2$ photon absorptions.

For large laser field intensities, which means large z arguments of the Bessel functions the following asymptotic expansion can be used for a fixed index [21]

$$J_n(z) = \sqrt{2/(\pi z)} Cos(z - n\pi/2 - \pi/4). \tag{17}$$

Which means an approximate $1/\sqrt{sin(\theta)cos(\theta)}$ angle dependence which and has a strong decay for large scattering angles. Note, that even this function shows very rapid oscillations. Figure 4 shows the same kind of cross sections for for $z = 10000$ (which means $I = 4.56 \times 10^{15}$ W/cm^2 intensity) and for $z = 6.61 \times 10^6$ (which means $I = 2.0 \times 10^{21}$ W/cm^2 intensity), respectively. Only the $n = 1$ one photon absorption process is considered.

IV. SUMMARY

We presented a formalism which gives analytic angular differential cross section model for laser assisted proton nucleon scattering on a Woods-Saxon optical potential where the nth-order photon absorption is taken into account simultaneously. As an example the proton - ^{12}Ca collision was investigated at moderate 49 MeV proton energies. The



FIG. 4: The calculated angular differential cross sections for $n = 0$ and photon absorption. The black thin solid line is for $I = 4.56 \times 10^{15}$ W/cm^2 intensity ($\mathcal{I} = 10000$). The dashed gray line is close to the relativistic threshold with $I = 2.0 \times 10^{21}$ W/cm^2 laser field intensity ($\mathcal{I} = 6.61 \times 10^6$).

laser intensities vary from from 10^7 W/cm^2 to 10^{21} W/cm^2. We found that at 10^7 W/cm^2 laser intensities the elastic Born total cross section is a factor of 90 times higher than the same process where a single photon is absorbed. At higher laser field intensities this ratio is even higher. We hope that this study will stimulate laser assisted nuclear scattering collisions which will be done in the Romanian ELI facility.

V. ACKNOWLEDGEMENT

We thank for Prof. Gyula Bencze for useful discussions and comments.

[1] Di Piazza A., Müller C., Hatsagortsyan K.Z. and Keitel C.H., Rev. Mod. Phys. **84**, 1171 (2012).
[2] Ehlotzky F., Jaron A. and Kaminsky J.Z., Phys. Rep. **297**, 63 (1998).
[3] Gunst J, Litvinov Y.A., Keitel L.C. and Pálffy A., Phys. Rev. Lett. **112**, 082501 (2014).
[4] Kornev A.S. and Zon B.A., Las. Phys. Lett. **4**, 588 (2007).
[5] Habs D. *et al.*, Eur. Phys. J. D, **55**, 279 (2009).
[6] Rudchik A.D. *el al.*, Eur. Phys. J. A, **44**, 221 (2010).
[7] Woods R.D. and Saxon D.S., Phys. Rev. **95**, 577 (1954).
[8] Abdul-Jalil I. and Jackson D.F., J. Phys. G: Nucl. Phys. **5**, 1699 (1979).
[9] von Geramb H.V., *Microscopic Optical Potentials*, Lecture Notes in Physics, Springer-Verlag 1979.
[10] Greiner W. and Mahrun J.A., *Nuclear Models*, Springer, ISBN 3-540-59180, 1996.
[11] Hlophe L. *et al.*, Phys. Rev. C **88**, 064608 (2013) and arXiv:1310.8334v1.
[12] Bunkin F.V. and Fjodorov M.V., ZsETF **49**, 1215 (1965).
[13] Bunkin F.V., Fjodorov M.V. and Kazakov A. E., Usp. Fiz. Nauk. **107**, 559 (1972).
[14] Faisal F.H.M. J., Phys. B. Atom. Mol. Phys. **6**, L312 (1972).
[15] Gontier Y. and Rahman N.K., Lett. al Nuovo Cim. **9**, 537 (1974).
[16] Bergou J., Acta. Phys. Hung. **40**, 55 (1976).
[17] Kroll N. M. and Watson K.M., Phys. Rev. A **8**, 804 (1973).
[18] Faisal F.H.M., *Theory of Multiphoton Processes*, Plenum Press, New York, 1987, ISBN 0-306-42317-0.
[19] http://www.physics.nist.gov/cgi-bin/cuu/Value?mpsme.
[20] Barna I.F., Apagyi B. and Scheid W., J. Phys. G. **26**, (2000) 323.
[21] Abramowitz M. and Stegun I., *Handbook of Mathematical Functions* Dover Publication., Inc. New York (1968) Chapter 9, Page 368. Eq. 9.2.1.